

# Future Needs in Climate Modeling: Aerosol-Cloud Interactions

Ann Fridlind

NASA Goddard Institute for Space Studies

ACE Science Working Group Meeting  
19 June 2008

# What are “the problems”?

- microphysics
  - warm rain formation (GCCN, turbulence effects)
  - ice formation mechanisms (convective, stratus, wave clouds)

# What are “the problems”?

- microphysics
  - warm rain formation (GCCN, turbulence effects)
  - ice formation mechanisms (convective, stratus, wave clouds)
- dynamics
  - microscale → macroscale
  - entrainment, self-organization, waves
  - interaction with microphysics

# What are “the problems”?

- microphysics
  - warm rain formation (GCCN, turbulence effects)
  - ice formation mechanisms (convective, stratus, wave clouds)
- dynamics
  - microscale → macroscale
  - entrainment, self-organization, waves
  - interaction with microphysics
- emergent quantities
  - precipitation
  - radiative fluxes
  - others



# What is required to solve “the problems”?

- intensive field experiments

# What is required to solve “the problems”?

- intensive field experiments
  - detailed case studies
  - cloud-scale modeling
  - “process studies”
  - goal: understand all important physical processes
  - example: GEWEX Cloud System Study (GCSS) program

# What is required to solve “the problems”?

- intensive field experiments
  - detailed case studies
  - cloud-scale modeling
  - “process studies”
  - goal: understand all important physical processes
  - example: GEWEX Cloud System Study (GCSS) program
- observational statistical quantification attempts not enough!

# What is required to solve “the problems”?

- intensive field experiments
  - detailed case studies
  - cloud-scale modeling
  - “process studies”
  - goal: understand all important physical processes
  - example: GEWEX Cloud System Study (GCSS) program
- observational statistical quantification attempts not enough!
  - disentangling microphysics and dynamics too complex
  - easily misleading (cause, effect?)
  - ship tracks only in shallow marine Sc (mesoscale response?)
  - models generally *required* for quantification
  - and for climate prediction (parameterization development)

# Role of satellite data sets

- part of “the solution”

# Role of satellite data sets

- part of “the solution”
  - global coverage
  - seasonal cycles and longer
  - provide constraints on GCMs (necessarily limited)
  - test GCMs performance for current climate

# Role of satellite data sets

- part of “the solution”
  - global coverage
  - seasonal cycles and longer
  - provide constraints on GCMs (necessarily limited)
  - test GCMs performance for current climate
- the future

# Role of satellite data sets

- part of “the solution”
  - global coverage
  - seasonal cycles and longer
  - provide constraints on GCMs (necessarily limited)
  - test GCMs performance for current climate
- the future
  - GCMs becoming more like CRMs (two-moment microphysics)
  - look to CRMs for data needs (subgrid scale, PDF approach)
  - goal: collect an ERBE-like data set
  - can only use what we know to guess what we need



# Role of satellite data sets

- part of “the solution”
  - global coverage
  - seasonal cycles and longer
  - provide constraints on GCMs (necessarily limited)
  - test GCMs performance for current climate
- the future
  - GCMs becoming more like CRMs (two-moment microphysics)
  - look to CRMs for data needs (subgrid scale, PDF approach)
  - goal: collect an ERBE-like data set
  - can only use what we know to guess what we need
- setting the stage

# Role of satellite data sets

- part of “the solution”
  - global coverage
  - seasonal cycles and longer
  - provide constraints on GCMs (necessarily limited)
  - test GCMs performance for current climate
- the future
  - GCMs becoming more like CRMs (two-moment microphysics)
  - look to CRMs for data needs (subgrid scale, PDF approach)
  - goal: collect an ERBE-like data set
  - can only use what we know to guess what we need
- setting the stage
  - seasonally ice-free Arctic, Greenland melting, sea level rise
  - tools for comparing costs (sea walls versus reactors)
  - aerosol indirect effect bar charts lose significance, magnitude?
  - aerosol effects = what prevented early detection?
  - precipitation and regional climate prediction gain importance
  - long-term goals: understand cloud physics, make GCMs work

# Some generalizations

- what do we know?

# Some generalizations

- what do we know?
  - aerosol-cloud coupling appears strongest in warm (low) clouds
  - aerosol variability is high vertically but low horizontally
  - cloud variability is high in both directions
  - aerosols + dynamics  $\leftrightarrow$  cloud properties
  - aerosol number size distribution dominates composition
  - cloud-base Sc precipitation  $\propto (LWP/N_d)^{7/4}$ ?

# Some generalizations

- what do we know?
  - aerosol-cloud coupling appears strongest in warm (low) clouds
  - aerosol variability is high vertically but low horizontally
  - cloud variability is high in both directions
  - aerosols + dynamics  $\leftrightarrow$  cloud properties
  - aerosol number size distribution dominates composition
  - cloud-base Sc precipitation  $\propto (LWP/N_d)^{7/4}$ ?
- what does this mean?

# Some generalizations

- what do we know?
  - aerosol-cloud coupling appears strongest in warm (low) clouds
  - aerosol variability is high vertically but low horizontally
  - cloud variability is high in both directions
  - aerosols + dynamics  $\leftrightarrow$  cloud properties
  - aerosol number size distribution dominates composition
  - cloud-base Sc precipitation  $\propto (LWP/N_d)^{7/4}$ ?
- what does this mean?
  - we need high vertical resolution to lowest altitudes
  - we need high horizontal resolution
  - “high resolution” = order 10-100 m
  - we need to aim for aerosol *number size distribution*
  - we need ground-based data (sub-cloud aerosols, dynamics)
  - we'd like to know  $LWP$ ,  $N_d$ , and precipitation to 0.1 mm/d

# ACE Mission

- important decisions
  - radar properties (scanning?, sensitivity?)
  - orbit height (low enough?)
  - lidar properties (multiple beams?)
  - polarimeter (resolution?)

# ACE Mission

- important decisions
  - radar properties (scanning?, sensitivity?)
  - orbit height (low enough?)
  - lidar properties (multiple beams?)
  - polarimeter (resolution?)
- some general issues
  - radar resolution crucial (orbit height)
  - single beam sufficient for wide homogeneous clouds
  - but scanning radar would boost statistics enormously
  - multiple-beam lidar offers similar advantages
  - single beam can miss horizontal structure
  - how will *dry* aerosol NSD be cornered?



# Decision-making process

- cost-benefit analysis (qualitative)

# Decision-making process

- cost-benefit analysis (qualitative)
- face-to-face group decision-making (scientists)
  - unfunded activity (necessarily limited)
  - quantitative studies not possible
  - some vested interests

# Decision-making process

- cost-benefit analysis (qualitative)
- face-to-face group decision-making (scientists)
  - unfunded activity (necessarily limited)
  - quantitative studies not possible
  - some vested interests
- face-to-face group decision-making (technologists)
  - scientists sparse (e.g., Instrument Incubator Program)
  - powerful tendency to favor virtuosic technology
  - latest technology  $\neq$  best science results

# Decision-making process

- cost-benefit analysis (qualitative)
- face-to-face group decision-making (scientists)
  - unfunded activity (necessarily limited)
  - quantitative studies not possible
  - some vested interests
- face-to-face group decision-making (technologists)
  - scientists sparse (e.g., Instrument Incubator Program)
  - powerful tendency to favor virtuosic technology
  - latest technology  $\neq$  best science results
- “simulator” studies (let's get together)
  - German HALO aircraft instruments
  - DOE ARM ground-based radar facility
  - ESA EarthCARE mission

# Simulator Studies

- an approach (for clouds)
  - simulate response of instruments to simulated clouds
  - size-resolved microphysics helpful
  - stick to field experiment case studies

# Simulator Studies





- an approach (for clouds)
  - simulate response of instruments to simulated clouds
  - size-resolved microphysics helpful
  - stick to field experiment case studies
- not a cure-all
  - still need clear scientific questions
  - still face trade-offs

# Simulator Studies

- an approach (for clouds)
  - simulate response of instruments to simulated clouds
  - size-resolved microphysics helpful
  - stick to field experiment case studies
- not a cure-all
  - still need clear scientific questions
  - still face trade-offs
- part of the future (and ACE?)
  - doesn't need to be expensive or time-consuming
  - basic technology in hand (e.g., Quickbeam)
  - same technology useful for later science

## Example 1: Tropical Warm Pool—International Cloud Experiment

# 2.8 GHz Radar (S-Band) Reflectivity + Doppler Velocity







Source: Data courtesy Christopher Williams (NOAA), DOE ARM data archive



## Example 2: Mixed-Phase Arctic Cloud Experiment





## 94-GHz Radar (MMCR) and Lidar (HSRL)



Source: Data courtesy DOE ARM and Ed Eloranta / U. Wisc.

## Example 2: Mixed-Phase Arctic Cloud Experiment


## 94-GHz Radar Reflectivity



Source: QuickBeam (<http://reef.atmos.colostate.edu/haynes/radarsim/>), Bastiaan van Dierenhoven / NASA GISS

## Example 2: Mixed-Phase Arctic Cloud Experiment





## 94-GHz Radar Doppler Velocity



Source: QuickBeam (<http://reef.atmos.colostate.edu/haynes/radarsim/>), Bastiaan van Dierenhoven / NASA GISS

## Example 2: Mixed-Phase Arctic Cloud Experiment





## Lidar Backscatter Cross-Section



Source: Bastiaan van Dierenhoven / NASA GISS

## Example 2: Mixed-Phase Arctic Cloud Experiment





## Lidar Circular Depolarization



Source: Bastiaan van Dierenhoven / NASA GISS

## Example 2: Mixed-Phase Arctic Cloud Experiment

# Lidar Circular Depolarization



Source: Bastiaan van Dierenhoven / NASA GISS

# Summary

- problems
  - microphysical
  - dynamical
  - microscale → cloud-system scale

# Summary

- problems
  - microphysical
  - dynamical
  - microscale → cloud-system scale
- solutions
  - focused field experiments (solve problems)
  - satellite data sets (make GCMs work)



# Summary

- problems
  - microphysical
  - dynamical
  - microscale → cloud-system scale
- solutions
  - focused field experiments (solve problems)
  - satellite data sets (make GCMs work)
- future
  - GCMs more like CRMs
  - integrated view of aerosols, clouds, precipitation
  - focus on regional-scale climate and precipitation
  - field-constrained CRM results for mission design